AMENDMENT TO THE CLAIMS

Please amend the claims as follows:

1-2 (canceled)

3. (original) An assay method for determining an aminoglycoside comprising:

combining a sample suspected of containing the aminoglycoside with an antibody specific for the aminoglycoside and with a reagent of the formula

wherein A is CH₂NH₂, CHCH₃NH₂, or CHCH₃NHCH₃; B is H or OH; D is H or OH; E is NH₂ or OH; G' is NH₂, NHCH₃, NH-T, or NCH₃-T; J is H or OH; L is H, CH₃, or OH; Q is H or CH₂OH; Y' is H, C(=O)CH(OH)CH₂CH₂NH₂, or T; T is a label; and T is present in only one of G' or Y'; the reagent comprising the analyte analog of the aminoglycoside and forming a detectable complex with the antibody; and

determining the presence or amount of the detectable complex as a measure of the aminoglycoside in the sample.

 (original) The assay method of claim 3 wherein the label is selected from the group consisting of enzymes, fluorescent compounds, luminescent compounds, radioactive isotopes, polymers, and microparticles. (original) An assay method according to claim 3 in which A is CH₂NH₂, B is H, D is H, E is NH₂, G' is NHCH₃, J is OH, L is CH₃, Q is H, and Y' is

- 6-9 (canceled)
- (original) A test kit for determining an aminoglycoside in a sample comprising in packaged combination an antibody specific for the aminoglycoside and a reagent of the formula

wherein A is CH₂NH₂, CHCH₃NH₂, or CHCH₃NHCH₃; B is H or OH; D is H or OH; E is NH₂ or OH; G' is NH₂, NHCH₃, NH-T, or NCH₃-T; J is H or OH; L is H, CH₃, or OH; Q is H or CH₂OH; Y' is H, C(=O)CH(OH)CH₂CH₂NH₂, or T; T is a label; and T is present in only one of G' or Y'.

(previously presented) A test kit according to claim 10 in which A is CH₂NH₂, B is H, D is H, E is NH₂, G' is NHCH₃, J is OH, L is CH₃, Q is H, and Y' is